

M-THEORY ON SPIN(7) MANIFOLDS AND THEIR F-THEORY DUALS

based on

1307.5858 with Bonetti, Grimm

1309.2287 with Bonetti, Grimm, Palti

Tom Pugh

OUTLINE

- Introduction
- Spin(7) Manifolds from Calabi-Yau Quotients
- M-theory on Spin(7) Manifolds with F-theory duals
- Conclusions



OUTLINE

- Introduction
- Spin(7) Manifolds from Calabi-Yau Quotients
- M-theory on Spin(7) Manifolds with F-theory duals
- Conclusions



INTRODUCTION

- The compactification of F-theory on Calabi-Yau fourfolds has been studied in great detail.
- However in 8 dimensions the largest special holonomy group is $\text{Spin}(7)$ and not $\text{SU}(4)$.
- We may therefore ask what features would be seen by compactifying on a $\text{Spin}(7)$ holonomy manifold.
- To approach this problem we consider the compactification of M-theory on a $\text{Spin}(7)$ holonomy manifold and then infer the F-theory dual.



OUTLINE

- Introduction
- Spin(7) Manifolds from Calabi-Yau Quotients
- M-theory on Spin(7) Manifolds with F-theory duals
- Conclusions



CALABI-YAU FOURFOLDS AND SPIN(7) HOLONOMY MANIFOLDS

- The decomposition of the chiral spinor of $Spin(8)$ under $SU(4)$ gives

$$\mathbf{8} \rightarrow \mathbf{1} \oplus \mathbf{1} \oplus \mathbf{6}$$

- From the two covariantly constant spinors we can build

$$J_{m\bar{n}} = i\eta_1 \gamma_{m\bar{n}} \eta_2 \quad \Omega_{mnr s} = (\eta_1 + i\eta_2) \gamma_{mnr s} (\eta_1 + i\eta_2)$$

- Alternatively, decomposing under $Spin(7)$ gives

$$\mathbf{8} \rightarrow \mathbf{1} \oplus \mathbf{7}$$

- From which we can build

$$\Phi_{mnr s} = \eta \gamma_{mnr s} \eta$$



SPIN(7) HOLONOMY MANIFOLDS FROM CALABI-YAU QUOTIENTS

- We will consider Spin(7) holonomy manifolds constructed using the method described by Joyce.
- We quotient the Calabi-Yau by an anti-holomorphic isometric involution

$$\sigma^2 = \mathbb{1} \quad \sigma^*(g) = g$$

$$\sigma^*(\Omega) = e^{2i\theta}\bar{\Omega} \quad \sigma^*(J) = -J$$

- The Cayley calibration of the Spin(7) manifold produced in this way is then given by

$$\Phi = \frac{1}{\nu^2} \left(\frac{1}{\|\Omega\|} \operatorname{Re}(e^{-i\theta}\Omega) + \frac{1}{8} J \wedge J \right)$$



OUTLINE

- Introduction
- Spin(7) Manifolds from Calabi-Yau Quotients
- M-theory on Spin(7) Manifolds with F-theory duals
- Conclusions




SPIN(7) QUOTIENTS OF ELLIPTICALLY FIBERED CALABI-YAU FOURFOLDS

- For the reduction to have an F-theory dual, the Calabi-Yau fourfold must be elliptically fibered

$$P \equiv y^2 - x^3 - f(u^i) x z^4 - g(u^i) z^6 = 0$$

- We then consider anti-holomorphic involutions which are compatible with this structure

$$\sigma : u^i \rightarrow \sigma_{ah}(u^i) \quad P = 0 \rightarrow P = 0$$



SPIN(7) EFFECTIVE ACTION

- By considering the action of σ we can see how harmonic forms on the Calabi-Yau fit within the Spin(7) cohomology

$$J = v^0 \omega_0 + v^\alpha \omega_\alpha + v^i \omega_i \quad \{\omega_{\alpha_+}\} = H_{\mathbf{21}}^2(Z_8, \mathbb{R})$$


$$\{\omega_0 \wedge J, \omega_{\alpha_-} \wedge J, \omega_i \wedge J\} \subset H_{\mathbf{1S}}^4(Z_8, \mathbb{R}) \oplus H_{\mathbf{35A}}^4(Z_8, \mathbb{R})$$

- The fields of the M-theory effective action can then be expanded in this basis
- $H_{\mathbf{21}}^2(Z_8, \mathbb{R})$ perturbations of C_3
- $H_{\mathbf{1S}}^4(Z_8, \mathbb{R}) \oplus H_{\mathbf{35A}}^4(Z_8, \mathbb{R})$ perturbations of g_{mn}
- The effective action is then invariant under N=1 supersymmetry in 3d



F-THEORY LIFT

- When lifting to F-theory the extra quotient which forms the Spin(7) manifold gives a quotient of the usual F-theory circle




- The circle quotient projects out the 4d fields such that the surviving modes form 3d N=1 multiplets



4D EFFECTS OF SPIN(7) QUOTIENTS

- The new anti-holomorphic quotient acts on the x^3 direction



- The effect of the quotient is then only seen in the F-theory picture at the fixed points.
- When the F-theory limit is taken and the circle becomes large, this quotient is then pushed away restoring 4d N=1 supersymmetry.



OUTLINE

- Introduction
- Spin(7) Manifolds from Calabi-Yau Quotients
- M-theory on Spin(7) Manifolds with F-theory duals
- Conclusions



FURTHER WORK


- Spin(7) Manifolds which are not quotients
- Spin(7) resolutions and the role of the extra modes
 - Extra 3d N=1 Multiplets
 - Confined to the fixed points on the F-theory circle
- Relationship to Hořava-Witten Constructions and Heterotic/M-theory



CONCLUSIONS

- The compactification of M-theory on a Spin(7) holonomy manifold can be carried out in a way that remains compatible with M-theory/F-theory duality.
- For the Calabi-Yau quotients analyzed the quotient also acts on the F-theory circle and the localized effects are washed out upon taking the large circle limit.
- Several intrinsically Spin(7) effects remain to be studied.
- Can the SUSY and Lorentz breaking scales be separated?



The left side of the slide features a decorative design consisting of several vertical stripes of varying widths and shades of light gray. Overlaid on these stripes are five solid dark blue circles of different sizes, arranged in a cluster that tapers towards the bottom.

THANK YOU