

Twisted Cohomotopy implies M5-brane anomaly cancellation

Hisham Sati*, Urs Schreiber†

September 24, 2021

Abstract

We highlight what seems to be a remaining subtlety in the argument for the cancellation of the total anomaly associated with the M5-brane in M-theory. Then we prove that this subtlety is resolved under the hypothesis that the C-field flux is charge-quantized in the generalized cohomology theory called J-twisted Cohomotopy.

Contents

1	Introduction	1
2	The issue	3
3	A resolution	6
3.1	For generic G_7 -Bianchi identity	6
3.2	Via Hypothesis H	8
A	Appendix: Tangent structure on sphere bundles	11

1 Introduction

Formulating M-theory remains an open problem (e.g. [Du96, 6][HLW98, p. 2][Du98, p. 6][NH98, p. 2][Du99, p. 330][Moo14, 12][CP18, p. 2][Wi19, @21:15][Du19, @17:04]). Even formulating just the field-theoretic decoupling limit of the worldvolume theory of M5-branes in M-theory remains an open problem (e.g. [La19, 6.3]). Nevertheless, it is traditionally assumed that enough is known about M-theory in general, and about M5-branes in particular, that it makes sense to check whether field theoretic anomalies (following [AW84][AG85]) on M5-brane worldvolumes cancel against M-theoretic anomaly inflow (following [CH85]) from the bulk spacetime (reviewed in the current context in [Ha05]).

Relevance of anomaly cancellation for M-theory. What from the physics perspective are called *anomalies* is what from the perspective of mathematics are *obstructions* (a point highlighted in [KS04][SSS09]). Hence such a cancellation of the total M5-brane anomaly, if properly identified, is strictly necessary for M-theory to exist: any remaining anomaly is an obstruction against the existence of the theory of which it is an anomaly. But conversely, wherever a putative anomaly in M-theory is found *not* to vanish, by available reasoning, this signifies (with the assumption that M-theory does in fact exist) the presence of a new aspect of the elusive

*hsati@nyu.edu

†us13@nyu.edu (corresponding author)

theory that had hitherto been missed: There must then be a new detail in the theory, previously unrecognized, which does imply the cancellation of the remaining anomaly, after all.

For this reason a careful mathematical analysis of anomaly cancellation in M-theory is in order. The tacit assumption that the proverbial magic of M-theory will take care of all cancellations anyway, freeing us from the burden of patient rigorous checks, would work only if the actual formulation of M-theory were known. Since it is not known, the situation is the reverse: A carefully deduced failure of anomalies to cancel provides a hint as to the actual formulation of the elusive theory.

Historical background on M5-brane anomaly cancellation. Indeed, the original computation of the total M5-brane anomaly in [Wi96b, 5] found the total anomaly *not* to vanish; and highlighted that the issue remains an open problem (“somewhat puzzling” [Wi96b, p. 35]). In reaction, several authors argued for several fixes, but, it seems, without convincing success (see [FHMM98, p.2] for pointers). Finally, [FHMM98, 3] argued that there is a previously neglected summand in the bulk anomaly inflow which needs to be taken into account (the top right term in diagram (5) below). That correction to the bulk anomaly inflow term has since become accepted (e.g., in [BBMN18, (5)]) as the solution to the M5-brane anomaly cancellation. The authors of [BBMN19a, A.4-5] recently recall the argument of [FHMM98] in streamlined form. Nonetheless, these arguments remain non-rigorous even by physics standards, due to a lack of actual formulation of M-theory. This is clearly acknowledged and highlighted by one of these authors, in [Ha05, p. 46].¹

Remaining issue. In this note we point out, in §2 below, that there does still remain one issue with the currently accepted anomaly cancellation argument [FHMM98, 3][BBMN19a, A.4-5] in itself. This is a simple observation: these authors made an *Ansatz* (see (6) below) for the C-field configuration ([FHMM98, (2.3)][BBMN19a, (A.18)]) which is seemingly not the most general admissible under the given assumptions (as also noticed in [Mon15, (3.12)][BBMN19b, (3.16)][BBM20, (2.34)]). Entering their anomaly cancellation argument instead with a general C-field configuration seemingly leaves one anomaly contribution uncanceled, shown on the bottom right of (5) below.

Resolution by Hypothesis H. We prove in §3 that this previously neglected remaining anomaly term does in fact vanish, hence that the anomaly cancellation argument of [Wi96b, §5][FHMM98, §3][BBMN19a, A.4-5] is completed, if one assumes a hypothesis [Sa13, §2.5] about the proper nature of the C-field in M-theory which in [FSS19b][FSS19c][SS19a][SS19b][SS21] we called *Hypothesis H*, recalled in §3.2 below. This hypothesis says that the M-theory C-field is charge-quantized (21) in the generalized cohomology theory called *J-twisted Cohomotopy* (24). We have previously demonstrated that this hypothesis implies a wealth of further anomaly cancellation conditions [FSS19b][FSS19c][SS19a][BSS19] and other effects [SS19b][SS21] expected in M-theory (exposition in [Sc20]).

Outlook. Since *Hypothesis H* gives rigorous mathematical meaning to the M-theoretic nature of the C-field, our derivation in §3 is a rigorous mathematical proof of the vanishing of the remaining anomaly term (5) from this hypothesis and, as such, completes the argument of [Wi96b, §5][FHMM98, §3][BBMN19a, A.4-5]. We do not claim to make the rest of that argument rigorous. In order to do so one will need also a rigorous definition of the M5-brane coupled to this C-field. We have presented results going towards that goal in [FSS19d][FSS20a][FSS20b][FSS20c][FSS19c], but more needs to be done.

Acknowledgement. We thank Domenico Fiorenza for very useful discussions. We thank an anonymous referee for alerting us of [BBMN19b], whose Section 4.1 overlaps with the discussion in our §3.1; see Remark 6.

¹[Ha05, p. 46]: “[...] the solution is not so clear. [The established procedure of anomaly cancellation] will not work for the M5-brane. [...] something new is required. What this something new is, is not a priori obvious. [...] [This is] a daunting task. To my knowledge no serious attempts have been made to study the problem. [...] [The proposal of [FHMM98]] probably should not be viewed as a final understanding of the problem. One would eventually hope for a microscopic formulation of M-theory which makes some of the manipulations [proposed in [FHMM98]] appear more natural.”

2 The issue

The geometry under consideration. We are dealing with (for background see e.g. [Du99][MiSc06]):

- (i) families of
- (ii) C-field configurations on
- (iii) 11-dimensional spacetimes
- (iv) sourced by magnetic 5-branes
- (v) of unit charge.

We now say what this means precisely: First, (i) with (iii) means that

$$X := \underbrace{X^{11}}_{\text{spacetime manifold}} \times \underbrace{U}_{\text{parameter manifold}}$$

is the product of an 11-dimensional manifold (spacetime) with a parameter manifold U of any dimension, while (ii) means that we consider a closed differential 4-form on X :

$$\underbrace{G_4}_{\text{family of C-field flux densities}} \in \Omega_{\text{cl}}^4(X) \implies \forall_{s \in U} \left(\underbrace{G_4^{(s)}}_{\text{C-field flux density at parameter } s} \in \Omega_{\text{cl}}^4(X^{11}) \right),$$

which hence is, in particular, a U -parametrized family of differential 4-forms on X^{11} .² Moreover, (iv) means, just as in Dirac's argument for magnetic 0-branes (e.g. [Al85, §2]), that X^{11} is the complement of a 5-brane worldvolume, hence that X is an orthogonal S^4 -fiber bundle (see Def. 13) as shown on the left of (1).

$$\begin{array}{ccc}
 \begin{array}{c} \text{unit sphere} \\ \text{around} \\ \text{M5-brane} \end{array} & S^4 \longrightarrow X & \begin{array}{c} \text{spacetime} \\ \text{(families)} \end{array} \\
 & \downarrow \text{4-sphere} \\
 & \text{fiber bundle} & S(p) \\
 \begin{array}{c} \text{parameter} \\ \text{manifold} \end{array} \times (0, \infty) \times \begin{array}{c} \text{M5-brane} \\ \text{worldvolume} \\ \text{(families)} \end{array} = Q & & \\
 \begin{array}{c} \text{radial} \\ \text{distance} \\ \text{from brane} \end{array} & & \\
 \end{array}
 \quad
 \begin{array}{ccc}
 \begin{array}{c} \text{C-field} \\ \text{4-flux density} \\ \text{(in families)} \end{array} & [G_4] \in & \begin{array}{c} \text{de Rham} \\ \text{cohomology} \end{array} & \begin{array}{c} \text{real} \\ \text{cohomology} \end{array} \\
 & \downarrow \text{total flux} \\
 & \text{through } S^4 & \downarrow \int_{S^4} & \downarrow \text{fiber integration} \\
 \begin{array}{c} \text{single M5} \\ (\Leftrightarrow \text{abelian 2-form field}) \end{array} & 1 \in & H_{\text{dR}}^{\bullet}(Q) & \simeq & H^{\bullet}(Q; \mathbb{R}) \\
 & & \downarrow S(p)_* & & \downarrow S(p)_*
 \end{array}
 \quad (1)$$

Finally, (v) means that the corresponding fiber integration (1) of G_4 over the 4-sphere fibers is unity³

$$S(p)_*[G_4] = 1 \in H^0(Q; \mathbb{R}) \quad (2)$$

as shown on the right of (1). This concludes the conditions (i)-(v) imposed on the spacetime geometry.

The general solution to (2) is the sum of half the Euler class of the S^4 -fibration (e.g. [BT82, §11][BC97, (2.3)]) with any *basic* class (by exactness of the Gysin sequence, e.g. [BT82, 14.33]), namely one pulled back from the base of the fibration:

²The inclined reader may think of the 4-flux data G_4 as being a value at stage U of the *mapping stack* $\mathbf{Fields}(X^{11}) := [X^{11}, \underline{\Omega}^4]$ into the sheaf of differential 4-forms, and of the anomaly polynomials (5) as being (classes of) differential forms on this mapping stack. While this is the correct point of view (exposition in [FSS13]), we will not further dwell on it here.

³Our derivations in §3 immediately apply generally to any integer charge $S(p)_*[G_4] \in \mathbb{N}$ (2). But for $N \geq 2$ even the nature of the higher gauge field on the M5-brane(s) remains open (see [FSS20a] for pointers and for a resolution for $N = 2$) and it seems premature to extrapolate the existing computations of worldvolume anomalies to this case (compare [HMM98, below (2.4)]).

$$\begin{array}{c}
\text{general} \\
\text{4-flux density} \\
\text{with unit flux} \\
\text{through } S^4 \\
[G_4] \\
= \\
\frac{1}{2} \mathcal{X}_4 + S(p)^* [G_4^{\text{basic}}] \\
\begin{array}{c}
\uparrow \\
\text{Euler class of} \\
S^4\text{-fibration} \\
\text{basic component:} \\
\text{pulled back from} \\
\text{base of } S^4\text{-fibration} \\
\uparrow \\
[G_4^{\text{basic}}] \\
\text{4-class on base} \\
\text{of } S^4\text{-fibration}
\end{array}
\end{array}
\quad
\begin{array}{c}
\in H^4(S^4) \\
\uparrow i_x^* \\
\in H^4(X) \\
\uparrow S(p)^* \\
\in H^4(Q)
\end{array}
\quad (3)$$

Remark 1 (The $1/2$ BPS M5 configuration and its generalization). The local model of the situation (1) is the trivial S^4 -fibration of the near horizon geometry of the smooth $1/2$ -BPS black M5-brane solution of 11-dimensional supergravity ([GT93], reviewed in [AFHS98, §2.1.2]), restricted to the Poincaré patch of 7-dimensional anti-de-Sitter spacetime:

$$\begin{array}{ccc}
S^4 & \longrightarrow & \text{AdS}_7^{\text{Poin}} \times S^4 \\
& & \downarrow S(p)=\text{pr}_1 \\
\mathbb{R}^{5,1} \times (0, \infty) & \underset{\text{diff}}{\simeq} & \text{AdS}_7^{\text{Poin}} \\
\begin{array}{c} \text{M5-brane} \\ \text{worldvolume} \end{array} & & \begin{array}{c} \text{radial} \\ \text{distance} \end{array} & & \begin{array}{c} \text{Poincaré chart of} \\ \text{anti-de Sitter spacetime} \end{array}
\end{array}
\quad G_4 = \text{vol}_{S^4} \quad (4)$$

So the point of (1) is to generalize the situation away from this highly symmetric $1/2$ -BPS configuration (4) to more general 5-brane configurations. While few to no black M5-brane solutions to 11d supergravity beyond (4) are known explicitly, only their topological structure matters for the discussion of anomaly cancellation; and that topological structure is (essentially by definition) what is expressed by (1).

Remark 2 (G_4 is singular on the M5-brane locus). Condition (2) implies (immediately so by the Poincaré Lemma, since G_4 is closed) that the flux density G_4 can *not* be extended to the locus of the M5-brane itself, which is (or would be) at the center $r = 0 \in [0, \infty)$ of the punctured ball $S^4 \times (0, \infty)$ in (1). Instead it must/would have a singularity at $r = 0$, as is manifest also from the basic example (4). Parts of the literature gloss over this subtlety; and the point made in [FHMM98, p. 4-5] was to argue that this is the source of the missing anomaly cancellation of [Wi96b]. To handle the singularity mathematically, these authors declared⁴ to multiply G_4 by a smooth radial cutoff function, thus rendering it no longer closed [FHMM98, (2.3), (3.4)] but, mathematically, extendable to the brane locus. Luckily, the key computation [FHMM98, (3.3)], recalled in (5) below, applies just as well if instead one leaves G_4 intact but removes the singular locus from spacetime, just as usual in supergravity (4).

Remark 3 (Focus on real cohomology). We focus here entirely on the anomaly polynomials in real cohomology, hence ignoring all torsion contributions (which become visible in integral cohomology) as well as all “global” anomaly contributions (which become visible in differential cohomology, see [FSS20d] for discussion of all these notions of cohomology and their relations). Because, while vanishing of the anomaly in real cohomology is not sufficient for full anomaly cancellation (which must happen in differential integral cohomology), it is the *necessary* first step. No argument about torsion of global contributions to the M5 anomaly (which, of course, one will eventually want to address) can affect the proof of anomaly cancellation at the rational/real approximation; and as long as subtleties do remain here, it behooves us to first focus on these. Therefore, we often abbreviate $H^\bullet(-) := H^\bullet(-; \mathbb{R})$, here and in the following.

⁴[FHMM98, p. 4]: “We leave to the future the very interesting question of the relation of this approach to that based on a direct study of solutions to supergravity.”

The anomaly polynomials. The cohomology classes contributing to the total M5-brane anomaly in the situation (1) are given in the literature as follows:

$$\begin{array}{c}
\text{Bulk} \\
\text{spacetime} \\
\text{CS-terms} \\
H^{12}(X) \\
\downarrow S(p)_* \text{ anomaly inflow} \\
H^8(Q) \\
\text{M5-brane} \\
\text{worldvolume} \\
\text{anomalies}
\end{array}
\quad
\begin{array}{c}
[G_4 \wedge I_8] \\
\downarrow \\
I_8 \\
+ \frac{-1}{24} p_2(NQ) \\
+ \frac{-1}{2} [G_4^{\text{basic}} \wedge G_4^{\text{basic}}] \\
\downarrow \\
\frac{-1}{2} [G_4^{\text{basic}} \wedge G_4^{\text{basic}}]
\end{array}
\quad
\begin{array}{c}
+ \\
\frac{-1}{6} [G_4 \wedge G_4 \wedge G_4] \\
\swarrow \quad \searrow \\
\frac{-1}{24} p_2(NQ) \quad \frac{-1}{2} [G_4^{\text{basic}} \wedge G_4^{\text{basic}}]
\end{array}
\quad
(5)$$

$$\begin{array}{c}
\underbrace{A_{\text{total}} = A_{\text{chiral fermion}} + A_{\text{chiral 2form}} + I_8 + \frac{-1}{24} p_2(NQ) + \frac{-1}{2} [G_4^{\text{basic}} \wedge G_4^{\text{basic}}]}_{\text{[Wi96b]: } \frac{1}{24} p_2(NQ)} \\
\underbrace{\hspace{10em}}_{\text{[FHMM98]: } 0} \\
\underbrace{\hspace{15em} + \frac{-1}{2} [G_4^{\text{basic}} \wedge G_4^{\text{basic}}]}_{\text{Hypothesis H: } 0}
\end{array}$$

We discuss the various items in (5):

- (i) The term I_8 (17) is the ‘‘one-loop polynomial’’ [DLM95][VW95], while the terms $A_{\text{chiral fermion}}$ and $A_{\text{chiral 2form}}$ are the plain anomalies [Wi96b, (5.1), (5.4)] of the chiral fermion and of the abelian chiral (i.e., with self-dual curvature) 2-form field in 6d QFT. These were expected in [Wi96b] to cancel against the influx of I_8 , but found there ([Wi96b, (5.7)]) to cancel only up to a remaining term $\frac{1}{24} p_2(NQ)$, where NQ denotes the normal bundle to the M5-brane locus in spacetime.
- (ii) The Chern-Simons term $-\frac{1}{6} G_4 \wedge G_4 \wedge G_4$ of 11-dimensional supergravity was argued in [FHMM98, §3] [BBMN19a, A.4-5] to contribute to the anomaly inflow from the bulk. Then a formula due to [BC97, Lem 2.1] shows that this gives rise to the previously missing summand of $\frac{-1}{24} p_2(NQ)$. However, these authors consider an Ansatz for the C-field configuration [FHMM98, (2.3), (3.4)][BBMN19a, (2.4)] which amounts to assuming
$$[G_4^{\text{basic}}] \stackrel{!}{=} 0 \quad (6)$$
in (3). If this assumption is not made, then the bulk Chern-Simons term in addition contributes an influx term $\frac{-1}{2} [G_4^{\text{basic}} \wedge G_4^{\text{basic}}]$ (bottom right of (5)), whose vanishing remains to be discussed.
- (iii) That the Ansatz (6) remained unjustified was pointed out in [FSS19a, (19)] and then in [BBMN19b, (3.16) & App. C] (where the basic component is denoted γ_4 , see also [BBM20, (2.34)]). Previously in [Mon15, (3.12)] the term G_4^{basic} was assumed to be non-vanishing, in general, and as a resolution it is was suggested [Mon15, (3.7)] that the traditional expression from [Wi96b, (5.7)] for the self-dual field anomaly $A_{\text{chiral 2form}}$ in real cohomology is wrong, in that it gets corrected by just the seemingly missing summand $\frac{\pm 1}{2} [G_4^{\text{basic}} \wedge G_4^{\text{basic}}]$. Unfortunately, we are unable to verify this derivation. Fortunately, it makes no difference:
- (iv) We prove in §3 that the seemingly restrictive Ansatz (6) is *implied* as soon as the dual G_7 -flux satisfies a Bianchi identity of a widely expected form (Theorem 4 below); in particular, if it satisfies the Bianchi identity that is implied by Hypothesis H (Thm. 9 below). In this way, Hypothesis H enforces the vanishing of the problematic remaining anomaly term by itself:

$$\text{Hypothesis H} \quad \Rightarrow \quad [G_4^{\text{basic}}] = 0 \quad \Rightarrow \quad [G_4^{\text{basic}} \wedge G_4^{\text{basic}}] = 0 \quad \text{in situation 1.}$$

This means, according to (5), that the total M5-brane anomaly is finally cancelled.

3 A resolution

We now prove that *Hypothesis H* implies, in the situation (1), the vanishing of the problematic basic term $[G_4^{\text{basic}}]$ in (3), thus implying the vanishing of the total M5-brane anomaly according to (5). We proceed in two steps:

- (1) In §3.1, we observe a general mechanism that applies as soon as the Bianchi identity $dG_7 = -\frac{1}{2}G_4 \wedge G_4$ holds with any correction by Pontrjagin classes (which is traditionally not guaranteed, see Rem. 7).
- (2) In §3.2, we discuss how Hypothesis H implements this mechanism.

Both steps rely on facts about tangent structure on sphere bundles, whose proofs we relegate to appendix A.

3.1 For generic G_7 -Bianchi identity

Theorem 4 (Vanishing of basic component). *Given a black M5-brane background (1) with C-field flux G_4 (3) satisfying a Bianchi identity of the form*

$$dG_7 = -\frac{1}{2}G_4 \wedge G_4 + P(p_1(\nabla^{TX}), p_2(\nabla^{TX})) \in \Omega_{\text{dR}}^8(X) \quad (7)$$

for P any polynomial of Pontrjagin forms, then the basic component of $[G_4]$ (3) vanishes:

$$[G_4^{\text{basic}}] = 0 \in H^4(B; \mathbb{R}). \quad (8)$$

Proof. The key point is that all Pontrjagin forms on a manifold that is an orthogonal spherical fibration are basic forms, by Prop. 22. This means with (7) that also the cup-square of the class of the 4-flux is basic:

$$[G_4]^2 = S(p)^*(P(p_1(NQ), p_2(NQ))) \in H^8(X; \mathbb{R}). \quad (9)$$

Consider then the fiber integration

$$S(p)_* : H^\bullet(X; \mathbb{R}) \longrightarrow H^{\bullet-4}(Q; \mathbb{R}) \quad (10)$$

along the fibers of $S(p)$, as in (1). By [BC97, Lemma 2.1], the fiber integration of the odd cup power χ_4^{2k+1} of the Euler class $\chi_4 \in H^4(X; \mathbb{R})$ of the fibration $S(p)$ are proportional to cup powers of the second Pontrjagin class of NQ :

$$S(p)_*(\chi_4^{2k+1}) = 2(p_2(NQ))^k \in H^{8k}(Q), \quad (11)$$

while the fiber integration of the even cup powers of the Euler class vanishes for all $k \in \mathbb{N}$:

$$S(p)_*(\chi_4^{2k}) = 0 \in H^{8k-4}(Q). \quad (12)$$

Notice also the *projection formula* (e.g. [BT82, Prop. 6.16][FSS18, (2)])

$$S(p)_*((S(p)^*\alpha) \wedge \beta) = \alpha \wedge S(p)_*\beta, \quad (13)$$

which in particular implies that the fiber integral of basic forms vanishes:

$$S(p)_*S(p)^*\alpha = S(p)_*(S(p)^*\alpha \wedge 1) = \alpha \wedge S(p)_*1 = 0. \quad (14)$$

Therefore, from (9) and by repeated use of formulas (11 - 14) we get:

$$\begin{aligned} 0 &= S(p)_*S(p)^*P(p_1(NQ), p_2(NQ)) \\ &= \frac{1}{2}S(p)_*[G_4 \wedge G_4] \\ &= \frac{1}{2}S(p)_*\left(\left(\frac{1}{2}\chi_4 + S(p)^*[G_4^{\text{basic}}]\right) \wedge \left(\frac{1}{2}\chi_4 + S(p)^*[G_4^{\text{basic}}]\right)\right) \\ &= \frac{1}{2}S(p)_*(\chi_4^2) + S(p)_*\left(\frac{1}{2}\chi_4 \wedge S(p)^*[G_4^{\text{basic}}]\right) + \frac{1}{2}S(p)_*(S(p)^*[G_4^{\text{basic}}])^2 \\ &= \frac{1}{2}S(p)_*(\chi_4^2) + S(p)_*\left(\frac{1}{2}\chi_4\right) \wedge [G_4^{\text{basic}}] + \frac{1}{2}S(p)_*S(p)^*[G_4^{\text{basic}} \wedge G_4^{\text{basic}}] \\ &= [G_4^{\text{basic}}]. \end{aligned} \quad (15)$$

□

Remark 5 (M5-Brane anomaly cancellation for traditional G_7 -Bianchi identity). A Bianchi identity for the (Hodge) dual flux density $G_7 := *_1 G_4$ of the form (7) is traditionally considered as ([DLM95, (1.1)][DFM03, (7.2)])

$$dG_7 = -\frac{1}{2}G_4^2 + I_8(p_1(\nabla^{TX}), p_2(\nabla^{TX})) \quad (16)$$

for the polynomial P being just the I_8 -term in (5):

$$I_8(p_1, p_2) := \frac{1}{48}(p_2 - \frac{1}{4}p_1^2). \quad (17)$$

Under this traditional assumption, Theorem 4 implies the condition (6) and hence the vanishing of the remaining M5-brane anomaly in (5).

Remark 6. Theorem 4 in conjunction with Remark 5 may be compared to an analogous physics argument in [BBMN19b, §4.1] (which appeared after [FSS19a] and during the writing of the first version of this article; we thank an anonymous referee for pointing this out).

Remark 7 (Further corrections to the G_7 -Bianchi identity). The traditional Bianchi identity (16) incorporates only one of several expected corrections to the plain supergravity Bianchi identity ($P = 0$). These M-theoretic *higher curvature corrections* are traditionally investigated via an action principle (e.g. [HT03][HO06][ST17]):

- (i) From the action principle one expects further higher derivative contributions to the Bianchi identity (16) ([ST17, (4.11)], following [HT03, around (56)]):

$$dG_7 = -\frac{1}{2}G_4^2 + I_8(p_1, p_2) + \overset{\text{further corrections}}{\delta\Delta L}, \quad (18)$$

which locally, on a chart $U \subset X$ where $G_4|_U = dC_3^U$ ([ST17, below (3.4)]), are of the form ([ST17, below (4.11)])

$$\delta\Delta L|_U = d\left(\frac{\delta}{\delta C_3^U} \cdots\right). \quad (19)$$

This shows at once that:

- (a) *locally on U* the correction is exact (which is the case highlighted in [ST17, below (4.11)]), but
 - (b) *globally on X* it fails to be exact as soon as G_4 is not globally exact (is only the curvature 4-form of a 2-gerbe connection with local connection 3-forms $\{C_3^U\}_{U \in \mathcal{U}}$ on an open cover \mathcal{U} of X , e.g. [FSS12, p. 22]), which is the generic case and the case of interest here, due to (2).
- (ii) The action principle, and hence any Bianchi identity derived from it, must moreover incorporate a global shift [Ts04, (4.16)]:

$$S(g, G_4) \mapsto S(g, G_4 + \frac{1}{4}p_1(\nabla^{TX})), \quad (20)$$

reflecting the expected [Wi96a, (1.2)][(1.2)][Wi96b] shifted flux quantization (23) of the C-field.

Apart from the general question of whether a classical action principle, of all things, can be the right principle for resolving foundations of M-theory, the complete form of the higher curvature correction in (18) remains open, and its combination with the shift (20) in the action principle seems not to have been discussed yet. Hence, under traditional assumptions, it remains unknown whether the assumption (7) is met in full M-theory.

What has been missing is a principle that fixes Bianchi identities on more fundamental grounds. Such a principle is *cohomological flux quantization* (21), to which we turn now in §3.2.

3.2 Via Hypothesis H

We briefly recall from [FSS19b][FSS20c] the motivation and formulation of *Hypothesis H* on the flux quantization principle for the C-field in M-theory. Then we show (Thm. 9 below) how this Hypothesis implements the above mechanism for cancellation of the remaining M5-anomaly term.

Flux/charge quantization of higher gauge fields. The key idea is that the mathematical nature of any higher gauge field is encoded in a *twisted generalized cohomology theory* $\tilde{A}^\tau(-)$, a notion known as *flux quantization* or *charge quantization* (see [Fr00][Sa10][FSS20d]): A generalized twisted *character map* [FSS20d, §5] approximates cocycles in τ -twisted A -cohomology by flux densities in twisted L_∞ -algebra valued de Rham cohomology, namely by differential forms satisfying polynomial differential relations – *Bianchi identities*:

$$\begin{array}{ccc}
 \begin{array}{c} \tau\text{-twisted } A\text{-cohomology} \\ \text{of spacetime } X \\ \tilde{A}^\tau(X) \end{array} & \xrightarrow{\text{twisted character map } \text{ch}_A^\tau} & \begin{array}{c} \tau_{\text{dR}}\text{-twisted} \\ L_\infty\text{-valued de Rham cohomology} \\ H_{\text{dR}}^{\tau_{\text{dR}}}(X; \mathcal{A}) \end{array} \\
 \begin{array}{c} \tau\text{-twisted } A\text{-cocycle} \\ [X \xrightarrow{c} A // G] \end{array} & \begin{array}{c} \text{rational approximation} \\ \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{\quad} \\ \text{flux quantization} \end{array} \end{array} & \begin{array}{c} \text{flux densities} \\ (F_i(c) \in \Omega_{\text{dR}}^\bullet(X))_{i \in I} \Big| \left(\begin{array}{c} \text{Bianchi identities} \\ dF_i(c) = P_i(\{F_j(c)\}_{j \in I}, \tau_{\text{dR}}) \end{array} \right)_{i \in I} \end{array} \\
 & & \begin{array}{c} \text{background} \\ \text{fluxes} \end{array}
 \end{array} \tag{21}$$

Flux/charge quantization in A -theory means to demand that the flux densities are in the image of the twisted A -character map (21) of an actual cocycle c in twisted A -cohomology, which then embodies the actual field configuration (its topological sector as shown here, for brevity, and the full field configuration after refinement to *differential* A -cohomology [FSS20d, §4.3]).

The archetypical example is Dirac’s flux quantization of the electromagnetic field (e.g. [Al85, §2][Fr00, §2]), which is the demand that the ordinary electromagnetic flux density F_2 (the Faraday tensor) is the character image of a cocycle in ordinary integral degree-2 cohomology $A(-) = H^2(-; \mathbb{Z})$ (hence is the curvature 2-form of a connection on a complex line bundle), which equivalently means that it represents an integral cohomology class $[F_2] \in H^2(X; \mathbb{Z}) \rightarrow H^2(X; \mathbb{R})$. Here the Bianchi identity obtained from (21) is the simple closure condition $dF_2 = 0$ ([FSS20d, Ex. 4.10]).

The most famous example is the K-theory conjecture in string theory [MM97][Wi98][Fr00] which states (review in [Wi00][Fr08]) that the B-field and the RR-field fluxes in type II string theory are jointly quantized in twisted [Wi98, §5.3][BM00] (and differential, see [GS19c] for recent rigorous developments) topological K-theory, $A^\tau(-) = \text{KU}^\tau(-)$. Indeed, the character map (21) in this case takes the following form ([FHT02, §2.5][FSS20d, Prop. 5.5], shown here for type IIA string theory, for definiteness):

$$\begin{array}{ccc}
 \begin{array}{c} \text{KU}^\tau(X) := \\ \tau\text{-twisted} \\ \text{complex K-theory} \\ \text{of } X \end{array} & \begin{array}{c} \left\{ \begin{array}{c} \mathbb{Z} \times \text{BU} // \text{BU}(1) \\ \downarrow \\ X \xrightarrow{\tau} \text{B}^2\text{U}(1) \\ \text{classifying map} \\ \text{of B-field} \end{array} \right\} \\ \text{classifying map} \\ \text{of RR-fields} \\ \text{vertical} \\ \text{homotopy} \end{array} & \xrightarrow{\text{twisted Chern character } \text{ch}_{\text{KU}}^\tau} & \left\{ \begin{array}{c} H_3, \quad \left. \begin{array}{l} dH_3 = 0, \quad [H_3] \in H^3(X; \mathbb{Z}) \\ \{F_{2k}\}_k \end{array} \right\} \\ \left. \begin{array}{l} dF_{2k+2} = H_3 \wedge F_{2k} \\ \text{RR-flux Bianchi identities} \end{array} \right\} / \sim \end{array} \tag{22}
 \end{array}$$

The Bianchi identities on the right of (22) are exactly those expected to be satisfied by the NS B-field flux H_3 and the RR-flux densities F_{2k} in type IIA string theory (see [FSS16b, §4][BMSS19, §1] for details and pointers).

But the M-theoretic lift of these H_3/F_{2k} -Bianchi identities (22) is (see [MaSa03, §4.2][Sa10][FSS16a, §3][BMSS19, §4]) just the G_7 -Bianchi identity (7) together with closure and the shifted integrality condition on the G_4 -flux:

$$dG_4 = 0, \quad \left[\overbrace{G_4 + \frac{1}{4}p_1(\nabla^{TX})}^{\tilde{G}_4} \right] \in H^4(X; \mathbb{Z}) \rightarrow H^4(X; \mathbb{R}). \tag{23}$$

Therefore, it is natural to ask for a cohomology theory whose twisted character map (21) enforces (7) and (23):

Hypothesis H is the statement [FSS19b, §2][FSS19c][SS19a][SS21] (following [Sa13, §2.5][FSS16a], review in [FSS19, §7]) that the cohomology theory for flux/charge quantization (21) of the C-field in M-theory is Borsuk-Spanier *Cohomotopy theory* $A^\tau(-) = \pi^\tau(-)$ [Sp49] in joint degrees 4 (for the M5-brane charge) and 7 (for the M2-brane charge) related by the quaternionic Hopf fibration and twisted by the tangent bundle via the J-homomorphism (“J-twist”):

The classifying space for degree- n Cohomotopy is (the homotopy type of) the n -sphere S^n , and for (orthogonally) twisted n -Cohomotopy it is the homotopy quotient (Borel construction) $S^n // O(n+1)$ of the canonical action of the orthogonal group $O(n+1)$ on $S^n \simeq S(\mathbb{R}^{n+1})$ (recalled as Def. 13 below):

$$\pi^\tau(X) := \left\{ \begin{array}{c} \text{cocycle} \\ \begin{array}{ccc} & S^4 // O(5) & \\ \swarrow c & \downarrow & \\ X & \xrightarrow{\tau} & BO(5) \end{array} \end{array} \right\} \Big/_{\text{vertical homotopy}} \xrightarrow[\text{[FSS20d, §5.3]}]{\text{twisted cohomotopical character } \text{ch}_\pi^\tau} \left\{ \begin{array}{l} G_4, \quad dG_4 = 0, \quad [G_4 + \frac{1}{4}p_1(\nabla^\tau)] \in H^4(X; \mathbb{Z}) \\ G_7, \quad dG_7 = -\frac{1}{2}G_4 \wedge G_4 + \frac{1}{8}p_2(\nabla^\tau) \end{array} \right\} /_{\sim} \quad (24)$$

On the right of (24) we are showing the form of the image of the character map (21) specified to orthogonally twisted 4-Cohomotopy (due to [FSS19b, Prop. 2.5 & 3.13], for more see [FSS20d, §5.3]). Both the $\frac{1}{4}p_1$ -shifted integral flux quantization on G_4 (23) is implied from charge-quantization in twisted Cohomotopy, as well as the general form of the G_7 -Bianchi identity (7). It just remains to relate the Pontrjagin classes of the twisting bundle τ to the tangent bundle:

The condition that the twist τ be compatible in degrees 7 and 4, along the quaternionic Hopf fibration $h_{\mathbb{H}}$ singles out ([FSS19b, Prop. 2.20]) the quaternionic central product subgroup $\text{Sp}(2) \cdot \text{Sp}(1) \subset \text{Spin}(8) \rightarrow \text{O}(8)$; and demanding that it, moreover, be compatible with factorization through the Atiyah-Penrose twistor fibration $t_{\mathbb{H}}$ (which corresponds [FSS20c][SS20c] to charge-quantization in *heterotic* M-theory) singles out ([FSS20c, Prop. 2.2]) the further subgroup $\text{Sp}(2) \subset \text{Sp}(2) \cdot \text{Sp}(1)$:

$$\begin{array}{ccc} \text{Sp}(2) \cdot \text{Sp}(1) & \xrightarrow{h_{\mathbb{H}}} & \text{Sp}(2) \cdot \text{Sp}(1) \\ \downarrow & \xrightarrow{\text{quaternionic Hopf fibration}} & \downarrow \\ S^7 & \xrightarrow{\quad} & S^4, \end{array} \quad \begin{array}{ccc} \text{Sp}(2) & \xrightarrow{h_{\mathbb{C}}} & \text{Sp}(2) \\ \downarrow & \xrightarrow{\text{complex Hopf fibration}} & \downarrow \\ S^7 & \xrightarrow{\quad} & \mathbb{C}P^3 \end{array} \quad \begin{array}{ccc} \text{Sp}(2) & \xrightarrow{t_{\mathbb{H}}} & \text{Sp}(2) \\ \downarrow & \xrightarrow{\text{Atiyah-Penrose twistor fibration}} & \downarrow \\ S^7 & \xrightarrow{\quad} & S^4. \end{array} \quad (25)$$

A key subtlety here is that the quaternionic unitary group $\text{Sp}(2)$ and the spin-group $\text{Spin}(5)$ are isomorphic as abstract Lie groups, but not as subgroups of $\text{Spin}(8) \rightarrow \text{O}(8)$ (nor are they conjugate subgroups); instead ([FSS19b, Prop. 2.17]) as subgroups they are mapped to each other under the triality automorphism on the ambient $\text{Spin}(8)$ -group:

$$\begin{array}{ccc} \text{Sp}(2) & \xrightarrow{\sim} & \text{Spin}(5) \\ \downarrow i_{\text{Sp}(2)} & \xrightarrow{\text{triality automorphism}} & \downarrow i_{\text{Spin}(5)} \\ \text{Spin}(8) & \xrightarrow{\text{tri}} & \text{Spin}(8), \end{array} \quad \begin{array}{ccc} B\text{Sp}(2) & \xrightarrow{\sim} & B\text{Spin}(5) \\ \downarrow Bi_{\text{Sp}(2)} & \xrightarrow{\text{delooped triality automorphism}} & \downarrow Bi_{\text{Spin}(5)} \\ B\text{Spin}(8) & \xrightarrow{B\text{tri}} & B\text{Spin}(8) \end{array} \quad (26)$$

$$\begin{array}{ccc} p_1 & \longleftrightarrow & p_1 \\ (\frac{1}{4}p_1)^2 - 24 \cdot I_8 & \longleftrightarrow & \frac{1}{4}p_2 \end{array}$$

As shown on the bottom right of (26), this triality automorphism, does not affect the first Pontrjagin class, but does induce a nontrivial transformation of Pontrjagin classes in degree 8 ([FSS19b, Lem. 2.19]). Therefore, as we consider tangential $\text{Sp}(2) \cdot \text{Sp}(1)$ -structure on spacetime to unify M2/M5-brane charge quantization in J-twisted Cohomotopy, and in fact tangential $\text{Sp}(2)$ -structure to account for charges in heterotic-theory, we arrive

A Appendix: Tangent structure on sphere bundles

Here we prove some results on (vertical) tangent structures to sphere bundles. The key consequences for the proofs in §3 are:

- (i) Prop. 22 (used in the proof of Thm.4 ,
- (ii) Prop. 19 (used in the proof of Thm. 9).

The first of these must be well-known to experts, but complete statements/proofs are hard to find in the literature (we give commented pointers to existing references as we proceed). We observe here that (i) follows as a direct corollary of the second statement (ii), which seems to be new. We give a slick homotopy-theoretic proof that neatly ties in with the formulation of *Hypothesis H*.

Homotopy theory. Following [FSS19b][FSS19c][SS21], we make free use of basic notions of homotopy theory (“higher structures”). For mathematical background and pointers see [FSS20d, §A][SS20b, §2]; for exposition in the context of string/M-theory see [JSSW19][FSS19]. This means that all topological spaces in the following are regarded up to weak homotopy equivalence (see [FSS20d, Ex. A.7]), which we denote by an equality sign, e.g. for $S^4 // \text{Spin}(5) = B\text{Spin}(4)$ in (38) below, where the double slash denotes the homotopy quotient or *Borel construction* for any topological/simplicial group G (see [NSS12b, Prop. 3.73])

$$X // G = H \times_G EG,$$

which subsumes the the classifying space construction $B(-)$ (e.g. [NSS12b, Ex. 3.68])

$$BG = * // G = * \times_G EG = (EG)/G$$

for *principal G -bundles*, being homotopy pullbacks (e.g. [FSS20d, A.24, A.27]) of the universal G -principal bundle

$$EG := G // G \rightarrow BG$$

([NSS12b, §4.1][SS20b, §2.2], traditional review in [Mit11]):

$$\begin{array}{ccc}
 P & \longrightarrow & G // G = EG \\
 \downarrow p & \text{homotopy pullback} & \downarrow \swarrow \\
 \text{principal } G\text{-bundle} & \text{(pb)} & \text{universal principal } G\text{-bundle} \\
 Q & \xrightarrow{\vdash P} & BG. \\
 & \text{classifying map} &
 \end{array} \tag{29}$$

Generally, for any (topological/simplicial) action $G \curvearrowright F$ of G on a typical fiber F , the homotopy quotient serves as the universal G -structured/associated F -fiber bundle ([NSS12a, §4][SS20b, §2.2]):

$$\begin{array}{ccc}
 E & \longrightarrow & F // G = EG \times_G F \\
 \downarrow p & \text{homotopy pullback} & \downarrow \swarrow \\
 \text{G-structured } F\text{-fiber bundle} & \text{(pb)} & \text{universal G-structured } F\text{-fiber bundle} \\
 Q & \xrightarrow{\vdash E} & BG. \\
 & \text{classifying map} &
 \end{array} \tag{30}$$

Here and in the following, we are notationally suppressing the homotopies filling all these diagrams.

G-Structures. Throughout, $n \in \mathbb{N}$ denotes any natural number. All (fiber-)vector spaces and, in particular, all (vertical) tangent spaces are assumed to be finite-dimensional.

Definition 11 (*G-Structure*). Given a topological group G and a homomorphism $\phi : G \rightarrow \text{GL}(d)$, we say that

- (i) G -structure on a real vector bundle $\mathcal{V} \xrightarrow{p} Q$ is a factorization of its classifying map $Q \xrightarrow{\vdash \mathcal{V}} \text{BO}(d)$ through $B\phi : BH \rightarrow B\text{GL}(d)$;
- (ii) G -structure on a real smooth manifold M^d is G -structure on its tangent vector bundle.

Remark 12 (Literature on G -structure). The notion of G -structures as an efficient tool for controlling the geometry of super-string compactifications is discussed in [Lo01][GMPW04][Koe10][Gal11][DDG14][Pr16][MPT17]. Beware that tradition in Cartan geometry insists that the homomorphism $G \xrightarrow{\phi} \text{GL}(d)$ be injective (e.g. [ČS09, p. 46]). Since this demand excludes common examples of “ G -structures” like Spin structures (but also String structures, etc.; and metaplectic structures, etc.) without being necessary for the part of the theory that is relevant here, we do not impose it. In algebraic topology this more general notion is known as $(BG, B\phi)$ -structures (see [Koc96, §1.4]) or as *tangential structures* [GMTW06, Sec. 5] (observing here that the canonical inclusion $\text{O}(d) \hookrightarrow \text{GL}(d)$ is the maximal compact subgroup, so that $\text{BO}(d) \rightarrow \text{BGL}(d)$ is a homotopy equivalence). See [SS20b, §4.2] for extensive discussion, comparison and further pointers.

Spherical fibrations.

Definition 13 (Orthogonal n -sphere fiber bundle).

(i) We say that an S^n -fiber bundle $p : X \rightarrow Q$ is *orthogonal* if it is equivalent to unit sphere bundle $S(p) : S(\mathcal{V}) \rightarrow Q$ inside a real vector bundle $p : \mathcal{V} \rightarrow Q$ (whose structure group may always be taken to be the orthogonal group).

(ii) This means equivalently that X is associated (30) via a classifying map $Q \xrightarrow{\vdash X} \text{BO}(n+1)$ to the classifying space for the orthogonal group, which fits into a homotopy-pullback diagram of the following form:

$$\begin{array}{ccc}
 X & \longrightarrow & S^n // \text{O}(n+1) \\
 \downarrow p & \text{homotopy pullback} & \downarrow \\
 \text{orthogonal } n\text{-sphere bundle} & \text{(pb)} & \text{universal orthogonal } n\text{-sphere bundle} \\
 Q & \xrightarrow{\vdash X} & \text{BO}(n+1), \\
 & \text{classifying map} &
 \end{array} \tag{31}$$

where on the top right we have the homotopy quotient (the Borel construction) of the n -sphere $S^n \simeq S(\mathbb{R}^{n+1})$ by its canonical action of the orthogonal group.

(iii) More generally, if a topological group G acts continuously on S^n , then we say that a G -associated n -sphere fiber bundle $X \rightarrow Q$ is one fitting into a homotopy-pullback diagram of this form:

$$\begin{array}{ccc}
 X & \longrightarrow & S^n // G \\
 \downarrow p & \text{homotopy pullback} & \downarrow \\
 G\text{-structured } n\text{-sphere bundle} & \text{(pb)} & \text{universal } G\text{-structured } n\text{-sphere bundle} \\
 Q & \xrightarrow{\vdash X} & BG, \\
 & \text{classifying map} &
 \end{array} \tag{32}$$

Example 14 (Universal orthogonal n -sphere fiber bundle). Denoting the canonical inclusion of orthogonal groups by

$$\begin{array}{ccc}
 \text{O}(n) & \hookrightarrow & \text{O}(n+1) \\
 A & \mapsto & \text{diag}(1, A),
 \end{array} \tag{33}$$

the universal example of orthogonal n -sphere bundles (Def. 13) is equivalent to the map Bi_n of classifying spaces induced from (33):

$$\begin{array}{ccc}
 S^n & \longrightarrow & S^n // \text{O}(n+1) = \text{BO}(n) \\
 \downarrow & \text{(pb)} & \downarrow \\
 * & \longrightarrow & \text{BO}(n+1) \xleftarrow{Bi_n}
 \end{array} \tag{34}$$

This example is classical, see for instance [BC97, p. 4]. But it is just a special case of a more general phenomenon that will be useful for our purpose:

Example 15 (Universal G -structured n -sphere fiber bundle). Let

$$H \hookrightarrow G \tag{35}$$

be an inclusion of topological groups. Then the homotopy quotient (Borel construction) of their coset space G/H by its canonical residual left G -action is equivalent to the homotopy type of the classifying space of H ([FSS19b, Lem. 2.7]):

$$\begin{array}{ccc} G/H & \rightarrow & (G/H)//G = BH \\ \downarrow & \text{(pb)} & \downarrow \swarrow^{Bi} \\ * & \longrightarrow & BG \end{array} \tag{36}$$

Therefore, when the coset space G/H is in fact an n -sphere equipped with G -action, which happens in the following cases ([FSS19b, Rem. 2.9, Prop. 2.23])

Generic				Exceptional			
H	\hookrightarrow	G	G/H	H	\hookrightarrow	G	G/H
$O(n)$	\subset	$O(n+1)$	S^n	$\text{Sp}(1) \times \text{Sp}(1)$	\subset	$\text{Sp}(2)$	S^4
$SO(n)$	\subset	$SO(n+1)$		$SU(3)$	\subset	G_2	S^6
$\text{Spin}(n)$	\subset	$\text{Spin}(n+1)$		G_2	\subset	$\text{Spin}(7)$	S^7
$SU(n)$	\subset	$SU(n+1)$	S^{2n-1}	$\text{Spin}(7)$	\subset	$\text{Spin}(9)$	S^{15}
$\text{Sp}(n)$	\subset	$\text{Sp}(n+1)$	S^{4n-1}				

(37)

then the universal G -associated n -sphere bundle is equivalently the classifying space of H ([FSS19b, Prop. 2.8]):

$$\begin{array}{ccc} S^n & \rightarrow & S^n//G = BH \\ \downarrow & \text{(pb)} & \downarrow \swarrow^{Bi} \\ * & \longrightarrow & BG \end{array} \tag{38}$$

Vertical tangent bundles to spherical fibrations. We now show how this universal homotopy-theoretic construction of sphere bundles knows everything about their vertical tangent bundles.

Proposition 16 (Classifying map of frame bundle to n -sphere). *Under the identification on the right of (34), the homotopy fiber inclusion $\text{fib}(Bi_n)$ of S^n into the universal orthogonal n -sphere fiber bundle (Example 14) is the classifying map $\vdash \text{Fr}(S^n)$ for the orthogonal frame bundle $\text{Fr}_O(S^n) \rightarrow S^n$ of the n -sphere:*

$$\begin{array}{ccc} S^n & \xrightarrow{\text{classifying map of orthogonal frame bundle}} & BO(n) \\ \downarrow & \vdash \text{Fr}_O(S^n) & \downarrow Bi_n \\ * & \longrightarrow & BO(n+1) \end{array} \tag{39}$$

Proof. The long homotopy fiber sequence of Bi_n (e.g. [NSS12a, Def. 2.26], following from the pasting law [NSS12a, Prop. 2.23]) shows that the homotopy fiber inclusion of Bi_n classifies an $O(n)$ -principal bundle over the n -sphere whose total space is $O(n+1)$ with $O(n)$ -action induced by the canonical inclusion i_n (33):

$$\begin{array}{ccccc} O(n) & \hookrightarrow & O(n+1) & \longrightarrow & * \\ \downarrow & \text{(pb)} & \downarrow & \text{(pb)} & \downarrow \\ * & \longrightarrow & S^n & \xrightarrow{\text{fib}(Bi_n)} & BO(n) \\ & & \downarrow & \text{(pb)} & \downarrow Bi_n \\ & & * & \longrightarrow & BO(n+1) \end{array}$$

Therefore, it is sufficient to observe that we have an isomorphism of $O(n)$ -principal bundles

$$\begin{array}{ccc} \mathcal{O}(n+1) & \xrightarrow{\sim} & \text{Fr}(S^n) \\ A & \mapsto & (A \cdot v_1, \dots, A \cdot v_n) \in \text{Fr}_{A \cdot v_0}(S^n), \end{array}$$

where $v_0, \dots, v_n \in \mathbb{R}^{n+1}$ are the canonical basis vectors and where on the right we regard $S^n = S(\mathbb{R}^{n+1})$ with the induced identification of $T_{A \cdot v_0} S^n \simeq (A \cdot v_0)^\perp \subset \mathbb{R}^{n+1}$. \square

More generally:

Proposition 17 (Classifying map of H -frame bundle of H -coset realization of n -sphere). *Given a coset-space realization of the n -sphere $S^n \simeq G/H$ (37) induced from a subgroup inclusion $H \xrightarrow{i} G$ (35) of compact Lie groups, then under the identification on the right of (38) the homotopy fiber inclusion $\text{fib}(Bi)$ of S^n into the universal G -associated n -sphere fiber bundle (Example 15) is a classifying map for the H -principal bundle on the n -sphere which exhibits its canonical H -structure (Def. 11):*

$$\begin{array}{ccc} S^n & \xrightarrow{\text{classifying map of } H\text{-frame bundle}} & BH \\ \downarrow & \text{Fr}_H(S^n) \text{ (pb)} & \downarrow Bi \\ * & \longrightarrow & BG \end{array} \quad (40)$$

Proof. As before, the long homotopy fiber sequence of Bi (e.g. [NSS12a, Def. 2.26], following from the pasting law [NSS12a, Prop. 2.23]) shows that the homotopy fiber inclusion of Bi classifies an H -principal bundle over the n -sphere whose total space is G with H -action induced by the given subgroup inclusion:

$$\begin{array}{ccccc} H & \xleftarrow{i} & G & \longrightarrow & * \\ \downarrow & & \downarrow & & \downarrow \\ * & \longrightarrow & S^n & \xrightarrow{\text{fib}(Bi)} & BH \\ & & \downarrow & & \downarrow Bi \\ & & * & \longrightarrow & BG \end{array}$$

Therefore, it is sufficient to observe that $G \rightarrow G/H$ is an H -frame bundle that exhibits H -structure (Def. 11) on the tangent bundle $T(G/H)$. This is basic fact of Cartan geometry, laid out for instance in [ČS09, p. 53]. \square

Example 18 (Canonical Spin structure on n -spheres). For the generic coset space realization of the n -sphere from (37), $S^n \simeq \text{Spin}(n+1)/\text{Spin}(n)$, Prop. 17 says that the homotopy fiber inclusion of the map of classifying spaces $B\text{Spin}(n) \rightarrow B\text{Spin}(n+1)$ classifies a $\text{Spin}(n)$ -principal bundle of the form $\text{Spin}(n+1) \rightarrow S^n$ and that this is a Spin structure (Def. 11) on the n -sphere. A traditional proof of this fact is spelled out in detail in [No15, Thm. A.6.6], see also [Gu88, §2.a].

Proposition 19 (H -Structure on vertical tangent bundle of G -associated sphere bundle). *Given a coset-space realization of the n -sphere $S^n \simeq G/H$ (37) induced from a Lie subgroup inclusion $H \xrightarrow{i} G$ (35), then for a G -associated S^n -fiber bundle $X \xrightarrow{S(p)} Q$ (32):*

- (i) *the vertical tangent bundle carries an H -structure (Def. 11);*
- (ii) *whose associated G -principal bundle is the pullback along $S(p)$ of that to which X is associated.*

Proof. By the classification (30) of fiber bundles, $S(p)$ sits in a homotopy pullback square as on the right of the following pasting diagram

$$\begin{array}{ccccc}
 & & \vdash T_{S(pq)}X & & \\
 & \searrow & \curvearrowright & \searrow & \\
 S_q^n & \longrightarrow & X & \xrightarrow{\vdash T_{S(p)}X} & S^n // G \cong BH \\
 \downarrow & \text{(pb)} & \downarrow S(p) & \text{(pb)} & \downarrow \\
 * & \xrightarrow{x} & Q & \xrightarrow{\vdash X} & BG.
 \end{array} \tag{41}$$

Notice that if the top map in the right square classifies H -structure on the vertical tangent bundle, as indicated by its label, then the homotopy-commutativity of the right square is equivalent to claim (ii). Hence it is sufficient now to prove that the top right map indeed classifies H -structure on the vertical tangent bundle; which then also proves claim (i).

To that end, consider any point $q \in Q$ and write S_q^n for the sphere fiber over this point, as shown by the homotopy pullback square on the left of (41). By the pasting law ([NSS12a, Prop. 2.23]) it follows that the full rectangle is a homotopy pullback. Therefore Prop. 17 says that the composite top map in (41) classifies H -structure on the tangent bundle of S_q^n . Since this true for all q , it follows that the H -principal bundle classified by $\vdash T_{S(p)}X$ restricts on each sphere fiber S_q^n to that sphere's tangent H -structure. But this is the defining property of (H -structure on) the vertical tangent bundle of X . \square

Corollary 20 (Once-stabilized vertical tangent bundle of orthogonal sphere bundle is basic). *The once-stabilized vertical tangent bundle to an orthogonal sphere bundle $S(p) : S(\mathcal{V}) \rightarrow Q$ (31) is isomorphic to the pullback of its underlying vector bundle:*

$$\underbrace{T_{S(p)}(S(\mathcal{V}))}_{\substack{\text{vertical} \\ \text{tangent bundle} \\ \text{orthogonal} \\ \text{sphere bundle}}} \times \mathbb{R} \underset{\text{one-step stabiliz.}}{\simeq} \underbrace{S(p)^*(\mathcal{V})}_{\text{pullback of associated vector bundle}}. \tag{42}$$

A traditional proof of this statement is indicated in [Go16, Prop. 1.1.9].

Proof. For the orthogonal subgroup inclusion $O(n) \xrightarrow{i_n} O(n+1)$ (33), Prop. 19 gives a homotopy pullback diagram (42) of this form:

$$\begin{array}{ccccc}
 & & \vdash T_{S(p)}S(\mathcal{V}) & & \\
 & \searrow & \curvearrowright & \searrow & \\
 S(\mathcal{V}) & \longrightarrow & S^n // O(n+1) & \xrightarrow{\cong} & BO(n) \\
 S(p) \downarrow & \text{(pb)} & \downarrow & \swarrow Bi_n & \\
 Q & \xrightarrow{\vdash \mathcal{V}} & BO(n+1) & &
 \end{array} \tag{43}$$

Noticing that postcomposition with Bi_n manifestly corresponds to one-step stabilization of an orthogonal vector bundle, the homotopy-commutativity of this square is exactly the claim to be proven. \square

The following is stated without proof as [CrEs03, Fact 3.1], apparently reading between the lines of [Mil56, p. 403].

Corollary 21 (Once-stabilized tangent bundle of orthogonal sphere bundle is basic).

If the base space Q is a smooth manifold, then the once-stabilized tangent bundle of the total space of an orthogonal sphere bundle $S(p) : S(\mathcal{V}) \rightarrow X$ (Def. 13) is isomorphic to the pullback along $S(p)$ of the Whitney sum of the tangent bundle of the base with the given vector bundle:

$$T(S(\mathcal{V})) \times \mathbb{R} \simeq_Q S(p)^*(TQ \oplus_Q \mathcal{V}). \tag{44}$$

Proof. Consider the following sequence of bundle isomorphisms over the base space Q :

$$\begin{aligned} T(\mathcal{S}(\mathcal{V})) \times \mathbb{R} &\simeq_Q (S(p)^*(TQ)) \oplus_Q \left(T_{S(p)}(\mathcal{S}(\mathcal{V})) \times \mathbb{R} \right) \\ &\simeq_Q (S(p)^*(TQ)) \oplus_Q (S(p)^*\mathcal{V}) \\ &\simeq_Q S(p)^*(TQ \oplus_Q \mathcal{V}). \end{aligned}$$

Here:

- (a) The first step is a splitting of the short exact sequence of vector bundles

$$0 \rightarrow T_{S(p)}\mathcal{S}(\mathcal{V}) \hookrightarrow T(\mathcal{S}(\mathcal{V})) \xrightarrow{dS(p)} S(p)^*(TQ) \rightarrow 0$$

that defines the vertical tangent bundle $T_{S(p)}\mathcal{S}(\mathcal{V})$, and which splits as a special case of the general splitting of short exact sequences of real vector bundles over paracompact Hausdorff base spaces, in particular over smooth manifolds, by forming orthogonal complements with respect to any choice of a continuous fiberwise inner product.

- (b) The second step is Corollary 20.

- (c) The last step is the distributivity of pullback over Whitney sum of vector bundles. \square

In conclusion :

Proposition 22 (Stable characteristic classes on sphere bundles are basic). *Given an orthogonal sphere-fiber bundle $\mathcal{S}(\mathcal{V})$,*

(i) *every stable characteristic class – hence every polynomial $P(p_1, p_2, \dots)$ of Pontrjagin classes p_i – of its vertical tangent bundle is basic, i.e.: pulled back from the base space Q :*

$$P(p_1, p_2, \dots)(T_{S(p)}\mathcal{S}(\mathcal{V})) = S(p)^* \underbrace{P(p_1, p_2, \dots)(\mathcal{V})}_{\in H^\bullet(Q)};$$

(ii) *and if the base space Q is a smooth manifold then then analogous statement holds for every stable class of the full tangent bundle*

$$P(p_1, p_2, \dots)(T\mathcal{S}(\mathcal{V})) = S(p)^* \underbrace{P(p_1, p_2, \dots)(TQ \oplus_Q \mathcal{V})}_{\in H^\bullet(Q)};$$

Proof. Since a stable characteristic class of a vector bundle, such as a Pontrjagin class, is one that can be pulled back from any direct sum of that vector bundle with a trivial vector bundle, the first claim follows by Cor. 20 and the second by Cor. 21. \square

Conflict of interest statement. On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

- [AFHS98] B. Acharya, J. Figueroa-O’Farrill, C. Hull, and B. Spence, *Branes at conical singularities and holography*, Adv. Theor. Math. Phys. **2** (1999), 1249-1286, [arXiv:hep-th/9808014].
- [Al85] O. Alvarez, *Topological quantization and cohomology*, Comm. Math. Phys. **100** 2 (1985), 279-309, [euclid:cmp/1103943448].
- [AG85] L. Alvarez-Gaumé and P. Ginsparg, *The structure of gauge and gravitational anomalies*, Ann. Phys. **161** (1985) 423-490, [doi:10.1016/0003-4916(85)90087-9].
- [AW84] L. Alvarez-Gaumé and E. Witten, *Gravitational Anomalies*, Nucl. Phys. **B234** (1984) 269-330, [doi:10.1016/0550-3213(84)90066-X].

- [BBMN18] I. Bah, F. Bonetti, R. Minasian, and E. Nardoni, *Class \mathcal{S} Anomalies from M-theory Inflow*, Phys. Rev. **D99** (2019), 086020, [arXiv:1812.04016].
- [BBMN19a] I. Bah, F. Bonetti, R. Minasian, and E. Nardoni, *Anomaly Inflow for M5-branes on Punctured Riemann Surfaces*, J. High Energy Phys. **2019** (2019), 123, [arXiv:1904.07250].
- [BBMN19b] I. Bah, F. Bonetti, R. Minasian, and E. Nardoni, *Anomalies of QFTs from M-theory and Holography*, J. High Energy Phys. **125** (2020), [arXiv:1910.04166].
- [BBM20] I. Bah, F. Bonetti, and R. Minasian, *Discrete and higher-form symmetries in SCFTs from wrapped M5-branes*, J. High Energy Phys. **196** (2021), [arXiv:2007.15003].
- [Sp49] E. Spanier, *Borsuk's Cohomotopy Groups*, Ann. Math. **50** (1949), 203-245, [jstor:1969362].
- [BM00] P. Bouwknegt and V. Mathai, *D-branes, B-fields and twisted K-theory*, J. High Energy Phys. **0003** (2000), 007, [arXiv:hep-th/0002023].
- [BC97] R. Bott and A. S. Cattaneo, *Integral Invariants of 3-Manifolds*, J. Diff. Geom. **48** (1998), 91-133, [arXiv:dg-ga/9710001].
- [BT82] R. Bott and L. Tu, *Differential Forms in Algebraic Topology*, Graduate Texts in Mathematics 82, Springer 1982, [doi:10.1007/978-1-4757-3951-0].
- [BMSS19] V. Braunack-Mayer, H. Sati and U. Schreiber, *Gauge enhancement of Super M-Branes via rational parameterized stable homotopy theory*, Comm. Math. Phys. **371** (2019), 197-265, [arXiv:1806.01115].
- [BSS19] S. Burton, H. Sati and U. Schreiber, *Lift of fractional D-brane charge to equivariant Cohomotopy theory*, J. Geom. Phys. **161** (2021) 104034, [arXiv:1812.09679].
- [CH85] C. Callan and J. Harvey, *Anomalies and Fermion Zero Modes on Strings and Domain Walls*, Nucl. Phys. **B250** (1985) 427-436, [doi:10.1016/0550-3213(85)90489-4].
- [ČS09] A. Čap and J. Slovák, *Parabolic Geometries I – Background and general theory*, Amer. Math. Soc., 2009, [ISBN:978-1-4704-1381-1].
- [CP18] S. M. Chester and E. Perlmutter, *M-Theory Reconstruction from (2,0) CFT and the Chiral Algebra Conjecture*, J. High Energy Phys. **2018** (2018) 116, [arXiv:1805.00892].
- [CrEs03] D. Crowley and C. Escher, *A classification of S^3 -bundles over S^4* , Diff. Geom. App. **18** 3 (2003) 363-380, [doi:10.1016/S0926-2245(03)00012-3], [arXiv:math/0004147].
- [DDG14] U. Danielsson, G. Dibitetto, and A. Guarino, *KK-monopoles and G-structures in M-theory/type IIA reductions*, J. High Energy Phys. **1502** (2015) 096, [arXiv:1411.0575].
- [DFM03] E. Diaconescu, D. S. Freed, and G. Moore, *The M-theory 3-form and E_8 gauge theory*, In: Elliptic Cohomology, 44-88, Cambridge University Press, 2007, [arXiv:hep-th/0312069].
- [Du96] M. Duff, *M-Theory (the Theory Formerly Known as Strings)*, Int. J. Mod. Phys. **A11** (1996), 5623-5642, [arXiv:hep-th/9608117].
- [Du98] M. Duff, *A Layman's Guide to M-theory*, Abdus Salam Memorial Meeting, Trieste, Italy, 19 - 22 Nov 1997, pp.184-213, [arXiv:hep-th/9805177].
- [Du99] M. Duff (ed.), *The World in Eleven Dimensions: Supergravity, Supermembranes and M-theory*, Institute of Physics Publishing, Bristol, 1999, [ISBN:9780750306720].
- [Du19] M. Duff, in: G. Farmelo, *The Universe Speaks in numbers*, interview 14, 2019, [grahamfarmelo.com/the-universe-speaks-in-numbers-interview-14]
- [DLM95] M. J. Duff, J. T. Liu, and R. Minasian, *Eleven Dimensional Origin of String/String Duality: A One Loop Test*, Nucl. Phys. **B452** (1995), 261-282, [arXiv:hep-th/9506126].
- [FSS12] D. Fiorenza, H. Sati and U. Schreiber, *Extended higher cup-product Chern-Simons theories*, J. Geom. Phys. **74** (2013), 130-163, [doi:10.1016/j.geomphys.2013.07.011], [arXiv:1207.5449].
- [FSS13] D. Fiorenza, H. Sati, U. Schreiber, *A higher stacky perspective on Chern-Simons theory*, In: D. Calaque et al. (eds.) *Mathematical Aspects of Quantum Field Theories*, Mathematical Physics Studies, Springer 2014 pp. 153-211, [doi:10.1007/978-3-319-09949-1], [arXiv:1301.2580].

- [FSS16a] D. Fiorenza, H. Sati, and U. Schreiber, *Rational sphere valued supercocycles in M-theory and type IIA string theory*, J. Geom. Phys. **114** (2017), 91-108, [doi:10.1016/j.geomphys.2016.11.024], [arXiv:1606.03206].
- [FSS16b] D. Fiorenza, H. Sati, and U. Schreiber, *T-Duality from super Lie n-algebra cocycles for super p-branes*, Adv. Theor. Math. Phys. **22** (2018), 1209–1270, [arXiv:1611.06536].
- [FSS18] D. Fiorenza, H. Sati, and U. Schreiber, *T-duality in rational homotopy theory via L_∞ -algebras*, Geometry, Topology and Mathematical Physics **1** (2018); special volume in tribute of Jim Stasheff and Dennis Sullivan, [arXiv:1712.00758].
- [FSS19] D. Fiorenza, H. Sati, and U. Schreiber, *The rational higher structure of M-theory*, Proc. LMS-EPSRC Durham Symposium *Higher Structures in M-Theory*, Aug. 2018, Fortsch. Phys., 2019, [doi:10.1002/prop.201910017], [arXiv:1903.02834].
- [FSS19a] D. Fiorenza, H. Sati, U. Schreiber, §2.5 & §4.5 in *Twisted Cohomotopy implies M-Theory anomaly cancellation*, preprint version 1, arXiv:1904.10207v1, (later split off from [FSS19b] into the present article).
- [FSS19b] D. Fiorenza, H. Sati, U. Schreiber, *Twisted Cohomotopy implies M-Theory anomaly cancellation on 8-manifolds*, Comm. Math. Phys. **377** (3) (2020) 1961-2025, [doi:10.1007/s00220-020-03707-2], [arXiv:1904.10207].
- [FSS19c] D. Fiorenza, H. Sati and U. Schreiber, *Twisted Cohomotopy implies M5 WZ term level quantization*, Comm. Math. Phys. **384** (2021) 403–432, [arXiv:1906.07417] [doi:10.1007/s00220-021-03951-0].
- [FSS19d] D. Fiorenza, H. Sati, and U. Schreiber, *Super-exceptional embedding construction of the M5-brane*, J. High Energy Phys. **2020**, 107 (2020), [arXiv:1908.00042].
- [FSS20a] D. Fiorenza, H. Sati and U. Schreiber, *Twisted Cohomotopy implies twisted String-structure on M5-branes*, J. Math. Phys. 2021 (in print), [arXiv:2002.11093].
- [FSS20b] D. Fiorenza, H. Sati, and U. Schreiber, *Super-exceptional M5-brane model: Emergence of SU(2)-flavor sector*, [arXiv:2006.00012].
- [FSS20c] D. Fiorenza, H. Sati, and U. Schreiber, *Twistorial Cohomotopy Implies Green-Schwarz anomaly cancellation*, [arXiv:2008.08544].
- [FSS20d] D. Fiorenza, H. Sati, and U. Schreiber, *The character map in (twisted differential) non-abelian cohomology*, [arXiv:2009.11909].
- [Fr08] S. Fredenhagen, *Physical Background to the K-Theory Classification of D-Branes*, chapter in: D. Husemoeller et al., *Basic Bundle Theory and K-Cohomology Invariants*, Springer 2008, [doi:10.1007/978-3-540-74956-1_1].
- [Fr00] D. Freed, *Dirac charge quantization and generalized differential cohomology*, Surveys in Differential Geometry, Int. Press, Somerville, MA, 2000, pp. 129-194, [doi:10.4310/SDG.2002.v7.n1.a6], [arXiv:hep-th/0011220].
- [FHMM98] D. Freed, J. Harvey, R. Minasian, and G. Moore, *Gravitational Anomaly Cancellation for M-Theory Fivebranes*, Adv. Theor. Math. Phys. **2** (1998), 601-618, [arXiv:hep-th/9803205].
- [FHT02] D. Freed, M. Hopkins, and C. Teleman, *Twisted equivariant K-theory with complex coefficients*, J. of Topology. **1** 1, [arXiv:math/0206257].
- [Gal1] J. Gaillard, *On G-structures in gauge/string duality*, 2011, [cronfa:42569].
- [GMTW06] S. Galatius, I. Madsen, U. Tillmann and M. Weiss, *The homotopy type of the cobordism category*, Acta Math. **202** 2 (2009), 195-239 [arXiv:math/0605249S].
- [GMPW04] J. Gauntlett, D. Martelli, S. Pakis, and D. Waldram, *G-Structures and Wrapped NS5-branes*, Comm. Math. Phys. 247 (2004) 421-445, [arXiv:hep-th/0205050].
- [GT93] G. Gibbons and P. Townsend, *Vacuum interpolation in supergravity via super p-branes*, Phys. Rev. Lett. **71** (1993), 3754, [arXiv:hep-th/9307049].
- [Go16] W. Gollinger, *Madsen-Tillmann-Weiss Spectra and a Signature Problem for Manifolds*, PhD Thesis, Münster, 2016, [ncatlab.org/nlab/files/GollingerMTWSpectra.pdf]

- [GS19c] D. Grady and H. Sati, *Ramond-Ramond fields and twisted differential K-theory*, [arXiv:1903.08843].
- [Gu88] S. Gutt, *Killing spinors on spheres and projective spaces*, in A. Trautman, G. Furlan (eds.) *Spinors in Geometry and Physics*, World Scientific, 1988, [doi:10.1142/9789814541510].
- [Ha05] J. Harvey, *TASI 2003 Lectures on Anomalies*, [spire:692082], [arXiv:hep-th/0509097].
- [HMM98] J. Harvey, R. Minasian and G. Moore, *Non-abelian Tensor-multiplet Anomalies*, J. High Energy Phys. **9809** (1998), 004, [arXiv:hep-th/9808060].
- [HLW98] P. S. Howe, N. D. Lambert, and P. C. West, *The Self-Dual String Soliton*, Nucl. Phys. **B515** (1998), 203-216, [arXiv:hep-th/9709014].
- [HT03] P. Howe and D. Tsimpis, *On higher-order corrections in M theory*, J. High Energy Phys. **0309** (2003) 038, [arXiv:hep-th/0305129].
- [HO06] Y. Hyakutake and S. Ogushi, *Higher Derivative Corrections to Eleven Dimensional Supergravity via Local Supersymmetry*, J. High Energy Phys. **0602** (2006), 068, [arXiv:hep-th/0601092].
- [JSSW19] B. Jurčo, C. Saemann, U. Schreiber, and M. Wolf, *Higher Structures in M-Theory*, Fortsch. Phys. **67** 8-9 (2019), [arXiv:1903.02807].
- [Koc96] S. Kochman, *Bordism, Stable Homotopy and Adams Spectral Sequences*, Fields Institute Monographs, Amer. Math. Soc., 1996, [cds:2264210].
- [Koe10] P. Koerber, *Lectures on Generalized Complex Geometry for Physicists*, Fortsch. Phys. **59** (2011), 169-242, [arXiv:1006.1536].
- [KS04] I. Kriz and H. Sati, *M Theory, Type IIA Superstrings, and Elliptic Cohomology*, Adv. Theor. Math. Phys. **8** (2004) 345-395, [arXiv:hep-th/0404013].
- [La19] N. Lambert, *Lessons from M2's and Hopes for M5's*, Proceedings of the LMS-EPSC Durham Symposium: *Higher Structures in M-Theory*, August 2018, Fortsch. Phys. **67** (2019), 1910011, [arXiv:1903.02825].
- [Lo01] J. Lott, *The Geometry of Supergravity Torsion Constraints*, [arXiv:math/0108125]; published as: *Torsion constraints in supergeometry*, Comm. Math. Phys. **133** (1990), 563-615, [doi:10.1007/BF02097010].
- [MaSa03] V. Mathai and H. Sati, *Some Relations between Twisted K-theory and E_8 Gauge Theory*, J. High Energy Phys. **0403** (2004), 016, [arXiv:hep-th/0312033].
- [MiSc06] A. Miemiec and I. Schnakenburg, *Basics of M-Theory*, Fortsch. Phys. **54** (2006), 5-72, [arXiv:hep-th/0509137].
- [Mil56] J. Milnor, *On manifolds homeomorphic to the 7-sphere*, Ann. Math. **64** 2 (1956), 399-405, [doi:10.1142/9789812836878_0001].
- [Mit11] S. A. Mitchell, *Notes on principal bundles and classifying spaces* (2011), [ncatlab.org/nlab/files/MitchellPrincipalBundles.pdf]
- [MM97] R. Minasian and G. Moore, *K-theory and Ramond-Ramond charge*, J. High Energy Phys. **9711** (1997), 002, [arXiv:hep-th/9710230].
- [MPT17] R. Minasian, D. Prins, and H. Triendl, *Supersymmetric branes and instantons on curved spaces*, J. High Energy Phys. **10** (2017), 159, [arXiv:1707.07002].
- [Mon15] S. Monnier, *Global gravitational anomaly cancellation for five-branes*, Adv. Theor. Math. Phys. **19** (2015), 701-724, [arXiv:1310.2250].
- [Moo14] G. Moore, *Physical Mathematics and the Future*, talk at Strings 2014, <http://www.physics.rutgers.edu/~gmoore/PhysicalMathematicsAndFuture.pdf>
- [NH98] H. Nicolai and R. Helling, *Supermembranes and M(atr)ix Theory*, In: M. Duff et. al. (eds.), *Nonperturbative aspects of strings, branes and supersymmetry*, World Scientific, Singapore, 1999, [arXiv:hep-th/9809103].
- [NSS12a] T. Nikolaus, U. Schreiber, and D. Stevenson, *Principal ∞ -bundles – General theory*, J. Homotopy Rel. Struc. **10** 4 (2015), 749-801, [arXiv:1207.0248].

- [NSS12b] T. Nikolaus, U. Schreiber, and D. Stevenson, *Principal ∞ -bundles – Presentations*, J. Homotopy Rel. Struc. **10**, 3 (2015), 565-622, [arXiv:1207.0249].
- [No15] N. Nowaczyk, *Dirac Eigenvalues of higher Multiplicity*, PhD thesis, Regensburg, 2015, [arXiv:1501.04045].
- [Pr16] D. Prins, *On flux vacua, $SU(n)$ -structures and generalised complex geometry*, PhD thesis, Lyon, 2015, [arXiv:1602.05415].
- [Sa10] H. Sati, *Geometric and topological structures related to M-branes*, in: *Superstrings, geometry, topology, and C^* -algebras*, 181–236, Proc. Sympos. Pure Math., 81, Amer. Math. Soc. (2010), [arXiv:1001.5020].
- [Sa13] H. Sati, *Framed M-branes, corners, and topological invariants*, J. Math. Phys. **59** (2018), 062304, [arXiv:1310.1060].
- [SS19a] H. Sati and U. Schreiber, *Equivariant Cohomotopy implies orientifold tadpole cancellation*, J. Geom. Phys. **156** 103775 (2020) [arXiv:1909.12277].
- [SS19b] H. Sati and U. Schreiber, *Differential Cohomotopy implies intersecting brane observables via configuration spaces and chord diagrams*, [arXiv:1912.10425].
- [SS20b] H. Sati and U. Schreiber, *Proper Orbifold Cohomology*, [arXiv:2008.01101].
- [SS20c] H. Sati and U. Schreiber, *The character map in equivariant twistorial Cohomotopy*, [arXiv:2011.06533].
- [SS21] H. Sati and U. Schreiber, *M/F-Theory as Mf-Theory*, [arXiv:2103.01877].
- [SSS09] H. Sati, U. Schreiber, and J. Stasheff, *Fivebrane structures*, Rev. Math. Phys. **21** 10 (2009), 1197-1240, [arXiv:0805.0564].
- [Sc20] U. Schreiber, *Microscopic brane physics from Cohomotopy theory*, talk at: H. Sati (org.), *M-Theory and Mathematics*, NYUAD Research Institute, January 27-30, 2020 [ncatlab.org/schreiber/files/Schreiber-MTheoryMathematics2020-v200126.pdf]
- [ST17] B. Souères and D. Tsimpis, *The action principle and the supersymmetrisation of Chern-Simons terms in eleven-dimensional supergravity*, Phys. Rev. D **95** (2017), 026013, [arXiv:1612.02021].
- [Ts04] D. Tsimpis, *11D supergravity at $\mathcal{O}(l^3)$* , J. High Energy Phys. **0410** (2004), 046, [arXiv:hep-th/0407271].
- [VW95] C. Vafa and E. Witten, *A One-Loop Test Of String Duality*, Nucl. Phys. **B447** (1995), 261-270, [arXiv:hep-th/9505053].
- [Wi96a] E. Witten, *On Flux Quantization In M-Theory And The Effective Action*, J. Geom. Phys. **22** (1997), 1-13, [arXiv:hep-th/9609122].
- [Wi96b] E. Witten, *Five-Brane Effective Action In M-Theory*, J. Geom. Phys. **22** (1997), 103-133, [arXiv:hep-th/9610234].
- [Wi98] E. Witten, *D-Branes And K-Theory*, J. High Energy Phys. **9812** (1998), 019, [arXiv:hep-th/9810188].
- [Wi00] E. Witten, *Overview Of K-Theory Applied To Strings*, Int. J. Mod. Phys. **A16** (2001), 693-706, [arXiv:hep-th/0007175].
- [Wi19] E. Witten, in: G. Farmelo, *The Universe Speaks in numbers*, interview 5, 2019, [grahamfarmelo.com/the-universe-speaks-in-numbers-interview-5].

Hisham Sati, *Mathematics, Division of Science, New York University Abu Dhabi, UAE.*
hsati@nyu.edu

Urs Schreiber, *Mathematics, Division of Science, New York University Abu Dhabi, UAE; on leave from Czech Academy of Science, Prague.*
us13@nyu.edu